

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 October 2002 (24.10.2002)

PCT

(10) International Publication Number
WO 02/083140 A1

- (51) International Patent Classification⁷: **A61K 31/5025**, C07D 487/04 (74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).
- (21) International Application Number: PCT/US02/10881 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 8 April 2002 (08.04.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/282,824 10 April 2001 (10.04.2001) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicants (*for all designated States except US*): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **MERCK SHARP & DOHME LIMITED** [GB/GB]; Hertford Road, Hoddenson, Hertfordshire EN11 9BU (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **BARNETT, Stanley, F.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **OWENS, Andrew, Pate** [GB/GB]; Hertford Road, Hoddesdon, Hertfordshire EN11 9BU (GB).
- Published:
— with international search report
— with amended claims
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 02/083140 A1

(54) Title: INHIBITORS OF AKT ACTIVITY

(57) Abstract: The present invention is directed to compounds comprising a triazolo[4,3-b]pyridazine moiety which inhibit the activity of Akt, a serine/threonine protein kinase. The invention is further directed to chemotherapeutic compositions containing the compounds of this invention and methods for treating cancer comprising administration of the compounds of the invention.

BEST AVAILABLE COPY

TITLE OF THE INVENTION

INHIBITORS OF AKT ACTIVITY

BACKGROUND OF THE INVENTION

5 The present invention relates to quinazoline containing compounds that are inhibitors of the activity of one or more of the isoforms of the serine/threonine kinase, Akt (also known as PKB). The present invention also relates to pharmaceutical compositions comprising such compounds and methods of using the instant compounds in the treatment of cancer.

10 Apoptosis (programmed cell death) plays essential roles in embryonic development and pathogenesis of various diseases, such as degenerative neuronal diseases, cardiovascular diseases and cancer. Recent work has led to the identification of various pro- and anti-apoptotic gene products that are involved in the regulation or execution of programmed cell death. Expression of anti-apoptotic
15 genes, such as Bcl2 or Bcl-x_L, inhibits apoptotic cell death induced by various stimuli. On the other hand, expression of pro-apoptotic genes, such as Bax or Bad, leads to programmed cell death (Aams et al. *Science*, 281:1322-1326 (1998)). The execution of programmed cell death is mediated by caspase -1 related proteinases, including caspase-3, caspase-7, caspase-8 and caspase-9 etc (Thornberry et al. *Science*,
20 281:1312-1316 (1998)).

 The phosphatidylinositol 3'-OH kinase (PI3K)/Akt/PKB pathway appears important for regulating cell survival/cell death (Kulik et al. *Mol. Cell. Biol.* 17:1595-1606 (1997); Franke et al, *Cell*, 88:435-437 (1997); Kauffmann-Zeh et al. *Nature* 385:544-548 (1997) Hemmings *Science*, 275:628-630 (1997); Dudek et al.,
25 *Science*, 275:661-665 (1997)). Survival factors, such as platelet derived growth factor (PDGF), nerve growth factor (NGF) and insulin-like growth factor-1 (IGF-1), promote cell survival under various conditions by inducing the activity of PI3K (Kulik et al. 1997, Hemmings 1997). Activated PI3K leads to the production of phosphatidylinositol (3,4,5)-triphosphate (PtdIns(3,4,5)-P3), which in turn binds to,
30 and promotes the activation of, the serine/threonine kinase Akt, which contains a pleckstrin homology (PH)-domain (Franke et al *Cell*, 81:727-736 (1995); Hemmings *Science*, 277:534 (1997); Downward, *Curr. Opin. Cell Biol.* 10:262-267 (1998), Alessi et al., *EMBO J.* 15: 6541-6551 (1996)). Specific inhibitors of PI3K or dominant negative Akt/PKB mutants abolish survival-promoting activities of these
35 growth factors or cytokines. It has been previously disclosed that inhibitors of PI3K

(LY294002 or wortmannin) blocked the activation of Akt/PKB by upstream kinases. In addition, introduction of constitutively active PI3K or Akt/PKB mutants promotes cell survival under conditions in which cells normally undergo apoptotic cell death (Kulik et al. 1997, Dudek et al. 1997).

5 Analysis of Akt levels in human tumors showed that Akt2 is overexpressed in a significant number of ovarian (J. Q. Cheung *et al. Proc. Natl. Acad. Sci. U.S.A.* 89:9267-9271(1992)) and pancreatic cancers (J. Q. Cheung *et al. Proc. Natl. Acad. Sci. U.S.A.* 93:3636-3641 (1996)). Similarly, Akt3 was found to be overexpressed in breast and prostate cancer cell lines (Nakatani et al. *J. Biol. Chem.* 10 274:21528-21532 (1999)).

 The tumor suppressor PTEN, a protein and lipid phosphatase that specifically removes the 3' phosphate of PtdIns(3,4,5)-P₃, is a negative regulator of the PI3K/Akt pathway (Li et al. *Science* 275:1943-1947 (1997), Stambolic et al. *Cell* 95:29-39 (1998), Sun et al. *Proc. Natl. Acad. Sci. U.S.A.* 96:6199-6204 (1999)). 15 Germline mutations of PTEN are responsible for human cancer syndromes such as Cowden disease (Liaw et al. *Nature Genetics* 16:64-67 (1997)). PTEN is deleted in a large percentage of human tumors and tumor cell lines without functional PTEN show elevated levels of activated Akt (Li et al. *supra*, Guldberg et al. *Cancer Research* 57:3660-3663 (1997), Risinger et al. *Cancer Research* 57:4736-4738 20 (1997)).

 These observations demonstrate that the PI3K/Akt pathway plays important roles for regulating cell survival or apoptosis in tumorigenesis.

 Three members of the Akt/PKB subfamily of second-messenger regulated serine/threonine protein kinases have been identified and termed 25 Akt1/PKB α , Akt2/PKB β , and Akt3/PKB γ respectively. The isoforms are homologous, particularly in regions encoding the catalytic domains. Akt/PKBs are activated by phosphorylation events occurring in response to PI3K signaling. PI3K phosphorylates membrane inositol phospholipids, generating the second messengers phosphatidylinositol 3,4,5-trisphosphate and phosphatidylinositol 3,4-bisphosphate, 30 which have been shown to bind to the PH domain of Akt/PKB. The current model of Akt/PKB activation proposes recruitment of the enzyme to the membrane by 3'-phosphorylated phosphoinositides, where phosphorylation of the regulatory sites

of Akt/PKB by the upstream kinases occurs (B.A. Hemmings, *Science* 275:628-630 (1997); B.A. Hemmings, *Science* 276:534 (1997); J. Downward, *Science* 279:673-674 (1998)).

Phosphorylation of Akt1/PKB α occurs on two regulatory sites, Thr³⁰⁸ in the catalytic domain activation loop and on Ser⁴⁷³ near the carboxy terminus (D. R. Alessi *et al.* *EMBO J.* 15:6541-6551 (1996) and R. Meier *et al.* *J. Biol. Chem.* 272:30491-30497 (1997)). Equivalent regulatory phosphorylation sites occur in Akt2/PKB β and Akt3/PKB γ . The upstream kinase, which phosphorylates Akt/PKB at the activation loop site has been cloned and termed 3'-phosphoinositide dependent protein kinase 1 (PDK1). PDK1 phosphorylates not only Akt/PKB, but also p70 ribosomal S6 kinase, p90RSK, serum and glucocorticoid-regulated kinase (SGK), and protein kinase C. The upstream kinase phosphorylating the regulatory site of Akt/PKB near the carboxy terminus has not been identified yet, but recent reports imply a role for the integrin-linked kinase (ILK-1), a serine/threonine protein kinase, or autophosphorylation.

Inhibition of Akt activation and activity can be achieved by inhibiting PI3K with inhibitors such as LY294002 and wortmannin. However, PI3K inhibition has the potential to indiscriminately affect not just all three Akt isozymes but also other PH domain-containing signaling molecules that are dependent on PtdIns(3,4,5)-P₃, such as the Tec family of tyrosine kinases. Furthermore, it has been disclosed that Akt can be activated by growth signals that are independent of PI3K.

Alternatively, Akt activity can be inhibited by blocking the activity of the upstream kinase PDK1. No specific PDK1 inhibitors have been disclosed. Again, inhibition of PDK1 would result in inhibition of multiple protein kinases whose activities depend on PDK1, such as atypical PKC isoforms, SGK, and S6 kinases (Williams *et al.* *Curr. Biol.* 10:439-448 (2000)).

It is an object of the instant invention to provide novel compounds that are inhibitors of Akt/PKB.

It is also an object of the present invention to provide pharmaceutical compositions that comprise.


It is also an object of the present invention to provide a method for treating cancer that comprises administering such inhibitors of Akt/PKB activity.

SUMMARY OF THE INVENTION

The instant invention provides for compounds that inhibit of Akt/PKB activity. In particular, the compounds disclosed selectively inhibit one or two of the Akt/PKB isoforms. The invention also provides for compositions comprising such inhibitory compounds and methods of inhibiting Akt/PKB activity by administering the compound to a patient in need of treatment of cancer.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of the instant invention are useful in the inhibition of the activity of the serine/threonine kinase Akt. In a first embodiment of this invention, the inhibitors of Akt activity are illustrated by the formula A:



wherein

R^1 represents phenyl, furyl, thienyl or pyridinyl, any of which groups may be optionally substituted with one, two or three substituents, independently selected from:

- a) halogen;
- b) C_{1-4} alkyl;
- c) C_{1-4} alkoxy;
- d) cyano;
- e) di(C_{1-4} alkyl)amino;
- f) hydroxy;

R^2 represents amino- C_{1-6} alkyl, C_{1-4} alkylamino-(C_{1-6})alkyl, di(C_{1-4} alkyl)amino-(C_{1-6})alkyl, hydroxy-(C_{1-6})alkyl or C_{1-4} alkoxy-(C_{1-6})alkyl, any of which groups may be optionally substituted;

R^3 represents hydrogen or C_{1-6} alkyl; and

R^4 independently represents hydrogen, C_{1-6} -alkyl, halogen, HO- or


C_{1-6} alkyl-O;

r is 0, 1 or 2;

or a pharmaceutically acceptable salt or stereoisomer thereof.

5

In another embodiment the inhibitors of the instant invention are illustrated by the formula A-I:



10

wherein

R^2 is as defined with reference to formula A above;

R^4 is selected from: C_{3-7} cycloalkyl and phenyl, any of which groups may be optionally substituted;

15

r is 0, 1 or 2;

s is 0, 1, 2 or 3; and

R^5 independently represents halogen, C_{1-4} alkyl or C_{1-6} alkoxy;

or the pharmaceutically acceptable salts thereof.

20

Specific compounds of the instant invention include:

2,2,*N,N*-tetramethyl-*N*-(3-phenyl-[1,2,4]triazolo[3,4-*a*]phthalazin-6-yl)-propane-1,3-diamine

25 *N*'-[3-(4-Methoxy-phenyl)-[1,2,4]triazolo[4,3-*a*]phthalazin-6-yl]-2,2,*N,N*-tetramethyl-propane-1,3-diamine

or a pharmaceutically acceptable salt thereof.

As used herein, the expression "C₁₋₆ alkyl" includes methyl and ethyl groups, and straight-chained or branched propyl, butyl, pentyl and hexyl groups. Particular alkyl groups are methyl, ethyl, *n*-propyl, isopropyl, *tert*-butyl and 2,2-dimethylpropyl. Derived expressions such as "C₁₋₆ alkoxy" are to be construed accordingly.

As used herein, the expression "C₁₋₄ alkyl" includes methyl and ethyl groups, and straight-chained or branched propyl and butyl groups. Particular alkyl groups are methyl, ethyl, *n*-propyl, isopropyl and *tert*-butyl. Derived expressions such as "C₁₋₄ alkoxy" are to be construed accordingly.

Typical C₃₋₇ cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The expression "C₃₋₇ cycloalkyl(C₁₋₆)alkyl" as used herein includes cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

Typical C₄₋₇ cycloalkenyl groups include cyclobutenyl, cyclopentenyl and cyclohexenyl.

Typical aryl groups include phenyl and naphthyl, preferably phenyl.

The expression "aryl(C₁₋₆)alkyl" as used herein includes benzyl, phenylethyl, phenylpropyl and naphthylmethyl.

The term "halogen" as used herein includes fluorine, chlorine, bromine and iodine, especially fluorine or chlorine.

In a particular embodiment, R² represents amino-C₁₋₆ alkyl, C₁₋₄ alkylamino-(C₁₋₆)alkyl or di(C₁₋₄ alkyl)amino-(C₁₋₆)alkyl. Representative values of R² include but are not limited to dimethylaminomethyl, aminoethyl, dimethylaminoethyl, diethylaminoethyl, 3-dimethylaminopropyl, 3-methylaminopropyl, 3-dimethylamino-2,2-dimethylpropyl and 3-dimethylamino-2-methylpropyl.

For use in medicine, the salts of the compounds of formula I will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid,

citric acid, tartaric acid, carbonic acid or phosphoric acid. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with
5 suitable organic ligands, e.g. quaternary ammonium salts.

The present invention includes within its scope prodrugs of the compounds of formula I above. In general, such prodrugs will be functional derivatives of the compounds of formula I which are readily convertible *in vivo* into the required compound of formula I. Conventional procedures for the selection and
10 preparation of suitable prodrug derivatives are described, for example, in *Design of Prodrugs*, ed. H. Bundgaard, Elsevier, 1985.

Where the compounds according to the invention have at least one asymmetric center, they may accordingly exist as enantiomers. Where the compounds according to the invention possess two or more asymmetric centers, they
15 may additionally exist as diastereoisomers. It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed within the scope of the present invention.

The compounds of the instant invention are inhibitors of the activity of Akt and are thus useful in the treatment of cancer, in particular cancers associated with irregularities in the activity of Akt and/or GSK3. Such cancers include, but are
20 not limited to ovarian, pancreatic and breast cancer.

In an embodiment of the invention, the instant compound is a selective inhibitor whose inhibitory efficacy is dependent on the PH domain. In this embodiment, the compound exhibits a decrease in *in vitro* inhibitory activity or no *in vitro*
25 inhibitory activity against truncated Akt proteins lacking the PH domain.

In another embodiment of the invention, the instant compound is a selective inhibitor whose inhibitory efficacy is dependent on the region of the proteins between the PH domain and the kinase domain. (See Konishi *et al. Biochem. and Biophys. Res. Comm.* 216: 526-534 (1995), Figure 2) That region will be referred to
30 as the hinge region. In this embodiment, the compound exhibits a decrease in *in vitro* inhibitory activity or no *in vitro* inhibitory activity against truncated Akt proteins lacking the PH domain and the hinge region.

Such an inhibitor that is dependent on either the PH domain, the hinge region or both provides a particular advantage since the PH domains and hinge
35 regions in the three Akt isoforms lack the sequence homology that is present in the

rest of the protein, particularly the homology found in the kinase domains (which comprise the catalytic domains and ATP-binding consensus sequences). It is therefore observed that certain inhibitor compounds, such as those described herein, are not only selective for one or two isoforms of Akt, but also are weak inhibitors or fail to inhibit other kinases, such as PKA and PKC, whose kinase domains share some sequence homology with the kinase domains of the Akt/PKB isoforms. Both PKA and PKC lack a PH domain.

In a further embodiment, the instant compound is selected from the group of a selective inhibitor of Akt 1, a selective inhibitor of Akt 2 and a selective inhibitor of both Akt 1 and Akt 2.

In another embodiment, the instant compound is selected from the group of a selective inhibitor of Akt 1, a selective inhibitor of Akt 2, a selective inhibitor of Akt3 and a selective inhibitor of two of the three Akt isoforms.

In another embodiment, the instant compound is a selective inhibitor of all three Akt isoforms, but is not an inhibitor of one, two or all of such Akt isoforms that have been modified to delete the PH domain, the hinge region or both the PH domain and the hinge region.

The present invention is further directed to a method of inhibiting Akt activity which comprises administering to a mammal in need thereof a pharmaceutically effective amount of the instant compound.

The compounds of this invention may be administered to mammals, preferably humans, either alone or, preferably, in combination with pharmaceutically acceptable carriers, excipients or diluents, in a pharmaceutical composition, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients

which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents; such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, microcrystalline cellulose, sodium crosscarmellose, corn starch, or alginic acid; binding agents, for example starch, gelatin, polyvinyl-pyrrolidone or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to mask the unpleasant taste of the drug or delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a water soluble taste masking material such as hydroxypropylmethyl-cellulose or hydroxypropyl-cellulose, or a time delay material such as ethyl cellulose, cellulose acetate butyrate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water soluble carrier such as polyethyleneglycol or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a

thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as butylated hydroxyanisole or alpha-tocopherol.

5 Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example
10 sweetening, flavoring and coloring agents, may also be present. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

 The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin
15 or mixtures of these. Suitable emulsifying agents may be naturally-occurring phosphatides, for example soy bean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening,
20 flavouring agents, preservatives and antioxidants.

 Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, flavoring and coloring agents and antioxidant.

 The pharmaceutical compositions may be in the form of a sterile
25 injectable aqueous solutions. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

 The sterile injectable preparation may also be a sterile injectable oil-in-water microemulsion where the active ingredient is dissolved in the oily phase. For example, the active ingredient may be first dissolved in a mixture of soybean oil and
30 lecithin. The oil solution then introduced into a water and glycerol mixture and processed to form a microemulsion.

 The injectable solutions or microemulsions may be introduced into a patient's blood-stream by local bolus injection. Alternatively, it may be advantageous to administer the solution or microemulsion in such a way as to maintain a constant

circulating concentration of the instant compound. In order to maintain such a constant concentration, a continuous intravenous delivery device may be utilized. An example of such a device is the Deltec CADD-PLUS™ model 5400 intravenous pump.

5 The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension for intramuscular and subcutaneous administration. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable
10 solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

15 Compounds of Formula A may also be administered in the form of a suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter, glycerinated
20 gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compound of Formula A are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

25 The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles and delivery devices, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than
30 intermittent throughout the dosage regimen.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specific amounts, as well as any product which results, directly or indirectly, from combination of the specific ingredients in the specified amounts.

The instant compounds may also be co-administered with other well known therapeutic agents that are selected for their particular usefulness against the condition that is being treated. For example, the instant compounds may be useful in combination with known anti-cancer and cytotoxic agents.

- 5 Similarly, the instant compounds may be useful in combination with agents that are effective in the treatment and prevention of neurofibromatosis, retinosis, polycystic kidney disease, infections of hepatitis delta and related viruses and fungal infections. The instant compositions may also be useful in combination with other inhibitors of parts of the signaling pathway that links cell surface
10 growth factor receptors to nuclear signals initiating cellular proliferation. Thus, the instant compounds may be utilized in combination with inhibitors of prenyl-protein transferase, including protein substrate competitive inhibitors of farnesyl-protein transferase, farnesyl pyrophosphate competitive inhibitors of the activity of farnesyl-protein transferase and/or inhibitors of geranylgeranyl-protein trans-
15 ferase. The instant compositions may also be co-administered with compounds that are selective inhibitors of geranylgeranyl protein transferase or selective inhibitors of farnesyl-protein transferase. The instant compositions may also be administered in combination with a compound that has Raf antagonist activity.

- The compounds of the instant invention may also be
20 co-administered with other well known cancer therapeutic agents that are selected for their particular usefulness against the condition that is being treated. Included in such combinations of therapeutic agents are combinations with an antineoplastic agent. It is also understood that the instant compositions and combinations may be used in conjunction with other methods of treating cancer and/or tumors,
25 including radiation therapy and surgery.

- Additionally, compositions of the instant invention may also be useful as radiation sensitizers. For instance, radiation therapy, including x-rays or gamma rays that are delivered from either an externally applied beam or by implantation of tiny radioactive sources, may be used in combination with the instant compounds to treat
30 cancer.

If formulated as a fixed dose, such combination products employ the combinations of this invention within the dosage range described below and the other pharmaceutically active agent(s) within its approved dosage range. Combinations of the instant invention may alternatively be used sequentially with known pharma-

ceutically acceptable agent(s) when a multiple combination formulation is inappropriate.

The instant compositions may also be useful in combination with an integrin antagonist for the treatment of cancer, as described in U.S. Serial No. 09/055,487, filed April 6, 1998, which is incorporated herein by reference.

As used herein the term an integrin antagonist refers to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to an integrin(s) that is involved in the regulation of angiogenesis, or in the growth and invasiveness of tumor cells. In particular, the term refers to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha v \beta 3$ integrin, which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha v \beta 5$ integrin, which antagonize, inhibit or counteract binding of a physiological ligand to both the $\alpha v \beta 3$ integrin and the $\alpha v \beta 5$ integrin, or which antagonize, inhibit or counteract the activity of the particular integrin(s) expressed on capillary endothelial cells. The term also refers to antagonists of the $\alpha v \beta 6$, $\alpha v \beta 8$, $\alpha 1 \beta 1$, $\alpha 2 \beta 1$, $\alpha 5 \beta 1$, $\alpha 6 \beta 1$ and $\alpha 6 \beta 4$ integrins. The term also refers to antagonists of any combination of $\alpha v \beta 3$, $\alpha v \beta 5$, $\alpha v \beta 6$, $\alpha v \beta 8$, $\alpha 1 \beta 1$, $\alpha 2 \beta 1$, $\alpha 5 \beta 1$, $\alpha 6 \beta 1$ and $\alpha 6 \beta 4$ integrins. The instant compounds may also be useful with other agents that inhibit angiogenesis and thereby inhibit the growth and invasiveness of tumor cells, including, but not limited to angiostatin and endostatin.

When a composition according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

In one exemplary application, a suitable amount of an inhibitor of one or two of the Akt/PKB isoforms is administered to a mammal undergoing treatment for cancer. Administration occurs in an amount of inhibitor of between about 0.1 mg/kg of body weight to about 60 mg/kg of body weight per day, preferably of between 0.5 mg/kg of body weight to about 40 mg/kg of body weight per day. A particular therapeutic dosage that comprises the instant composition includes from about 0.01mg to about 1000mg of inhibitor of one or two of the Akt/PKB isoforms. Preferably, the dosage comprises from about 1mg to about 1000mg of inhibitor of one or two of the Akt/PKB isoforms.

Examples of an antineoplastic agent include, in general, microtubule-stabilising agents (such as paclitaxel (also known as Taxol®), docetaxel (also

known as Taxotere®), or their derivatives); alkylating agents, anti-metabolites; epidophyllotoxin; an antineoplastic enzyme; a topoisomerase inhibitor; procarbazine; mitoxantrone; platinum coordination complexes; biological response modifiers and growth inhibitors; hormonal/anti-hormonal therapeutic agents and haematopoietic growth factors.

Example classes of antineoplastic agents include, for example, the anthracycline family of drugs, the vinca drugs, the mitomycins, the bleomycins, the cytotoxic nucleosides, the taxanes, the epothilones, discodermolide, the pteridine family of drugs, diylenes and the podophyllotoxins. Particularly useful members of those classes include, for example, doxorubicin, carminomycin, daunorubicin, aminopterin, methotrexate, methopterin, dichloro-methotrexate, mitomycin C, porfiromycin, 5-fluorouracil, 6-mercaptopurine, gemcitabine, cytosine arabinoside, podophyllotoxin or podo-phyllotoxin derivatives such as etoposide, etoposide phosphate or teniposide, melphalan, vinblastine, vincristine, leurosine, vindesine, leurosine, paclitaxel and the like. Other useful antineoplastic agents include estramustine, cisplatin, carboplatin, cyclophosphamide, bleomycin, gemcitabine, ifosamide, melphalan, hexamethyl melamine, thiotepa, cytarabin, idatrexate, trimetrexate, dacarbazine, L-asparaginase, camptothecin, CPT-11, topotecan, ara-C, bicalutamide, flutamide, leuprolide, pyridobenzoindole derivatives, interferons and interleukins.

All patents, publications and pending patent applications identified are hereby incorporated by reference.

Abbreviations used in the description of the chemistry and in the Examples that follow are:

Ac ₂ O	Acetic anhydride;
Boc	t-Butoxycarbonyl;
DBU	1,8-diazabicyclo[5.4.0]undec-7-ene;


Reactions used to generate the compounds of this invention are prepared by employing reactions as shown in the Scheme 1, in addition to other standard manipulations such as ester hydrolysis, cleavage of protecting groups, etc., as may be known in the literature or exemplified in the experimental procedures. Substituents R, R^a and R^b, as shown in the Scheme, represent the substituents R¹ and

R²; however their point of attachment to the ring is illustrative only and is not meant to be limiting.

These reactions may be employed in a linear sequence to provide the compounds of the invention or they may be used to synthesize fragments which are subsequently joined by the alkylation reactions described in the Schemes.

SYNOPSIS OF SCHEME 1:

The requisite intermediates are in some cases commercially available, or can be prepared according to literature procedures. Thus as shown in Scheme 1, a suitably substituted 1,4-dichlorophthalazine is reacted with a stoichiometric amount of hydrazine to provide the 1-chloro-4-hydrazinophthalazine I, which can then undergo acylation/cyclization with an appropriate acid chloride to provide the triazolophthalazine II. The intermediate II can then react with a suitable amine to provide the instant compound III.

Reaction Scheme 1

EXAMPLES

Examples provided are intended to assist in a further understanding of the invention. Particular materials employed, species and conditions are intended to be further illustrative of the invention and not limitative of the reasonable scope thereof.

EXAMPLE 1

10 2,2,N,N-tetramethyl-N-(3-phenyl-[1,2,4]triazolo[3,4-a]phthalazin-6-yl)-propane-1,3-diamine (Compound 1)

Step 1: 1-Chloro-4-hydrazinophthalazine hydrochloride

To a stirred solution of hydrazine hydrate (40ml) in ethanol (120mL) at 80°C was added 1,4-dichlorophthalazine (20g). This reaction mixture was stirred at 80°C for 0.5 hours, then left to cool and the product was collected by filtration and dried under vacuum to give 1-chloro-4-hydrazinophthalazine hydrochloride (14.6g).
15 ¹H NMR (250 MHz, DMSO) • 4.64 (2H, vbs), 7.2 (1H, vbs), 7.92 (4H, bm).

20 Step 2: 6-Chloro-3-phenyl-1,2,4-triazolo[3,4-a]phthalazine

To a solution of 1-chloro-4-hydrazinophthalazine hydrochloride (10g) in dioxan (220ml) was added triethylamine (7.24ml) and benzoyl chloride (6.04ml). This mixture was heated at reflux for 8 hours under nitrogen. After cooling the reaction mixture was concentrated under vacuum and the solid obtained was collected by filtration, washed with water and diethyl ether and dried under vacuum, to yield
25 the title compound (12.0g). ¹H NMR (250 MHz, DMSO) • 7.60 (3H, m), 8.00 (1H, t, J=8.4Hz), 8.19 (1H, t, J=8.4Hz), 8.31 (3H, m), 8.61 (1H, d, J=6.3Hz).

30 Step 3: 2,2,N,N-tetramethyl-N-(3-phenyl-[1,2,4]triazolo[3,4-a]phthalazin-6-yl)-propane-1,3-diamine

The title compound was prepared as described in Example 1, Step 3, but replacing the 6-Chloro-7-cyclobutyl-3-phenyl-[1,2,4]triazolo[4,3-b]pyridazine with the 6-Chloro-3-phenyl-1,2,4-triazolo[3,4-a]phthalazine from Step 2.

¹H NMR (360MHz, CDCl₃) δ 1.13 (6H, s), 2.35 (2H, s), 2.46-2.50 (8H, m), 3.47 (2H, vbs), 7.16-7.27 (2H, m), 7.44-7.86 (5H, m), 8.55-8.57 (2H, m), 8.68 (1H, m). MS (ES+) MH⁺ = 375

EXAMPLE 2

N'-[3-(4-Methoxy-phenyl)-[1,2,4]triazolo[4,3-*a*]phthalazin-6-yl]-2,2,*N,N*-tetramethylpropane-1,3-diamine (Compound 2)

The title compound was prepared in an analogous fashion to Example 1, except substituting 3-fluorobenzoic hydrazine for the benzoic hydrazine in Step 2. ¹H NMR (360MHz, CDCl₃) δ 1.13 (6H, s), 2.45 (6H, s), 2.49 (2H, s), 3.45-3.46 (2H, m), 3.90 (3H, s) 7.04-7.07 (2H, m), 7.65-7.70 (2H, m), 7.80-7.84 (1H, m), 8.51 (2H, m), 8.66 (1H, m). MS (ES+) MH⁺ = 405

EXAMPLE 3

Cloning of the human Akt isoforms and ΔPH-Akt1

The pS2neo vector (deposited in the ATCC on April 3, 2001 as ATCC) was prepared as follows: The pRmHA3 vector (prepared as described in *Nucl. Acid Res.* 16:1043-1061 (1988)) was cut with BglII and a 2734 bp fragment was isolated. The pUChsneo vector (prepared as described in *EMBO J.* 4:167-171 (1985)) was also cut with BglII and a 4029 bp band was isolated. These two isolated fragments were ligated together to generate a vector termed pS2neo-1. This plasmid contains a polylinker between a metallothionine promoter and an alcohol dehydrogenase poly A addition site. It also has a neo resistance gene driven by a heat shock promoter. The pS2neo-1 vector was cut with Psp5II and BsiWI. Two complementary oligonucleotides were synthesized and then annealed (CTGCGGCCGC (SEQ.ID.NO.: 1) and GTACGCGGCCGCAG (SEQ.ID.NO.: 2)). The cut pS2neo-1 and the annealed oligonucleotides were ligated together to generate a second vector, pS2neo. Added in this conversion was a NotI site to aid in the linearization prior to transfection into S2 cells.

Human Akt1 gene was amplified by PCR (Clontech) out of a human spleen cDNA (Clontech) using the 5' primer:

5' CGCGAATTCAGATCTACCASTEAGCGACGTGGCTATTGTG 3'

(SEQ.ID.NO.: 3), and the 3' primer:

5'CGCTCTAGAGGATCCTCAGGCCGTGCTGCTGGC3' (SEQ.ID.NO.: 4). The 5'

primer included an EcoRI and BglII site. The 3' primer included an XbaI and BamHI

5 site for cloning purposes. The resultant PCR product was subcloned into pGEM3Z

(Promega) as an EcoRI / Xba I fragment. For expression / purification purposes, a

middle T tag was added to the 5' end of the full length Akt1 gene using the PCR

primer: 5'GTACGATGCTGAACGATATCTTCG 3' (SEQ.ID.NO.: 5). The resulting

PCR product encompassed a 5' KpnI site and a 3' BamHI site which were used to

10 subclone the fragment in frame with a biotin tag containing insect cell expression

vector, pS2neo.

For the expression of a pleckstrin homology domain (PH) deleted (Δ aa

4-129, which includes deletion of a portion of the Akt1 hinge region) version of Akt1.

PCR deletion mutagenesis was done using the full length Akt1 gene in the pS2neo

15 vector as template. The PCR was carried out in 2 steps using overlapping internal

primers (5'GAATACATGCCGATGGAAAGCGACAGGGGCTGAAGAGATGGA

GGTG 3' (SEQ.ID.NO.: 6), and 5'CCCCTCCATCTCTTCAGCCCCAGTCGC

TTTCCATCGGCATGTATTC 3' (SEQ.ID.NO.: 7)) which encompassed the deletion

and 5' and 3' flanking primers which encompassed the KpnI site and middle T tag on

20 the 5' end. The final PCR product was digested with KpnI and SmaI and ligated into

the pS2neo full length Akt1 KpnI / Sma I cut vector, effectively replacing the 5' end

of the clone with the deleted version.

Human Akt3 gene was amplified by PCR of adult brain cDNA

(Clontech) using the amino terminal oligo primer:

25 5' GAATTCAGATCTACCATGAGCGATGTTACCATTGTG 3' (SEQ.ID.NO.: 8);

and the carboxy terminal oligo primer:

5' TCTAGATCTTATTCTCGTCCACTTGCAGAG 3' (SEQ.ID.NO.: 9).

These primers included a 5' EcoRI / BglII site and a 3' XbaI / BglII site for cloning

purposes. The resultant PCR product was cloned into the EcoRI and XbaI sites of

30 pGEM4Z (Promega). For expression / purification purposes, a middle T tag was

added to the 5' end of the full length Akt3 clone using the PCR primer: 5'

GGTACCATGGAATACATGCCGATGGAAAGCGATGTTACCATTGTGAAG

3' (SEQ.ID.NO.: 10). The resultant PCR product encompassed a 5' KpnI site which

allowed in frame cloning with the biotin tag containing insect cell expression vector,

35 pS2neo.

Human Akt2 gene was amplified by PCR from human thymus cDNA (Clontech) using the amino terminal oligo primer:

5' AAGCTTAGATCTACCATGAATGAGGTGTCTGTC 3' (SEQ.ID.NO.: 11); and the carboxy terminal oligo primer: 5'

5 GAATTCGGATCCTCACTCGCGGATGCTGGC 3' (SEQ.ID.NO.: 12). These primers included a 5' HindIII / BglII site and a 3' EcoRI / BamHI site for cloning purposes. The resultant PCR product was subcloned into the HindIII / EcoRI sites of pGem3Z (Promega). For expression / purification purposes, a middle T tag was added to the 5' end of the full length Akt2 using the PCR primer: 5'

10 GGTACCATGGAATACATGCCGATGGAAAATGAGGTGTCTGTCATCAAAG 3' (SEQ.ID.NO.: 13). The resultant PCR product was subcloned into the pS2neo vector as described above.

EXAMPLE 4

15

Expression of human Akt isoforms and Δ PH-Akt1

The DNA containing the cloned Akt1, Akt2, Akt3 and Δ PH-Akt1 genes in the pS2neo expression vector was purified and used to transfect *Drosophila* S2 cells (ATCC) by the calcium phosphate method. Pools of antibiotic (G418, 500 μ g/ml) resistant cells were selected. Cell were expanded to a 1.0L volume ($\sim 7.0 \times 10^6$ / ml), biotin and CuSO₄ were added to a final concentration of 50 μ M and 50 mM respectively. Cells were grown for 72h at 27°C and harvested by centrifugation. The cell paste was frozen at -70°C until needed.

25

EXAMPLE 5

Purification of human Akt isoforms and Δ PH-Akt1

Cell paste from one liter of S2 cells, described in Example 13, was lysed by sonication with 50mls 1% CHAPS in buffer A: (50mM Tris pH 7.4, 1mM EDTA, 1mM EGTA, 0.2mM AEBSF, 10 μ g/ml benzamidine, 5 μ g/ml of leupeptin, aprotinin and pepstatin each, 10% glycerol and 1mM DTT). The soluble fraction was purified on a Protein G Sepharose fast flow (Pharmacia) column loaded with 9mg/ml anti-middle T monoclonal antibody and eluted with 75 μ M EYMPME (SEQ.ID.NO.: 14) peptide in buffer A containing 25% glycerol. Akt/PKB containing fractions were pooled and the protein purity evaluated by SDS-PAGE. The purified protein was

quantitated using a standard Bradford protocol. Purified protein was flash frozen on liquid nitrogen and stored at -70°C.

EXAMPLE 6

Kinase Assays

This procedure describes a kinase assay which measures phosphorylation of a biotinylated GSK3-derived peptide by human recombinant active Akt/PBK isoforms or Akt/PBK mutants. The ³³P-labeled biotinylated product can be captured and detected using Streptavidin coated Flashplates (NEN LifeSciences) or Streptavidin Membrane Filter Plates (Promega). Alternatively, a GSK3-derived peptide with 2 added lysine residues was used as the substrate and subsequently captured using Phosphocellulose Membrane Filter Plates (Polyfiltronics).

Materials:

Active human Akt: The following active human Akt isoforms were utilized in the *in vitro* assays: active human Akt1 (obtained from Upstate Biotechnology, catalog no. 14-276, 15 µg/ 37 µl (6.76 µM)) or recombinant lipid activated Akt1 (prepared as described in Example 5); Akt2 (prepared as described in Example 5); Akt3 (prepared as described in Example 5); and delta PH-Akt1 (prepared as described in Example 5).

Akt specific peptide substrate: GSK3α (S21) Peptide #3928, biotin-GGRARTSSFAEPG (SEQ.ID.NO.: 15), FW = 1517.8 (obtained from Macromolecular Resources) for Streptavidin Flashplate or Streptavidin Filter Plate detection.

GSK3α (S21) Peptide #G80613, KKGGRARTSSFAEPG (SEQ.ID.NO.: 14), FW = 1547.8 (obtained from Research Genetics) for Phosphocellulose filter plate detection.

Standard Assay Solutions:

A. 10X Assay Buffer: 500 mM HEPES, pH 7.5
1% PEG
1 mM EDTA

1 mM EGTA

20 mM β -Glycerol phosphate

- 5 B. Active Akt (500 nM): Diluent (1X Assay buffer, 10% glycerol, 0.1% β -mercaptoethanol, 1.0 μ M microcystin LR and 1.0 mM EDTA) was added to a vial containing 37 μ l of active Akt isoform (6.76 μ M). Aliquots were flash frozen in liquid N_2 and stored at $-70^\circ C$.
- 10 C. 1 mM Akt specific peptide substrate in 50 mM Tris pH 7.5, 1 mM DTT.
- D. 100 mM DTT in di H_2O .
- 15 E. 100X Protease Inhibitor Cocktail (PIC): 1 mg/ml benzamidine, 0.5 mg/ml pepstatin, 0.5 mg/ml leupeptin, 0.5 mg/ml aprótinin.
- F. 3 mM ATP, 200 mM $MgCl_2$ in H_2O , pH 7.9.
- 20 G. 50% (v/v) Glycerol.
- H. 1% (wt/v) BSA (10 mg/ml) in di H_2O , 0.02% (w/v) NaN_3 .
- I. 125 mM EDTA.
- 25 J. 0.75% (wt/v) Phosphoric Acid.
- K. 2.5 M Potassium Chloride.
- 30 L. Tris Buffered Saline (TBS), 25 mM Tris, 0.15 M Sodium Chloride, pH 7.2 (BupH Tris Buffered Saline Pack, Pierce catalog no. 28376).

Procedure for Streptavidin Flash Plate Assay:Step 1:

5 A 1 μ l solution of the test compound in 100% DMSO was added to 20 μ l of 2X substrate solution (20 μ M GSK3 Peptide, 300 μ M ATP, 20 mM MgCl_2 , 20 μ Ci / ml [$\gamma^{33}\text{P}$] ATP, 1X Assay Buffer, 5% glycerol, 1 mM DTT, 1X PIC, 0.1% BSA and 100 mM KCl). Phosphorylation reactions were initiated by adding 19 μ l of 2X Enzyme solution (6.4 nM active Akt/PKB, 1X Assay Buffer, 5% glycerol, 1 mM DTT, 1X PIC and 0.1% BSA). The reactions were then incubated at room temperature for 45
10 minutes.

Step 2:

The reaction was stopped by adding 170 μ l of 125 mM EDTA. 200 μ l of stopped reaction was transferred to a Streptavidin Flashplate[®] PLUS (NEN Life Sciences, catalog no. SMP103). The plate was incubated for ≥ 10 minutes at room temperature
15 on a plate shaker. The contents of each well was aspirated, and the wells rinsed 2 times with 200 μ l TBS per well. The wells were then washed 3 times for 5 minutes with 200 μ l TBS per well with the plates incubated at room temperature on a platform shaker during wash steps.

20 The plates were covered with sealing tape and counted using the Packard TopCount with the appropriate settings for counting [^{33}P] in Flashplates.

Procedure for Streptavidin Filter Plate Assay:Step 1:

25 The enzymatic reactions as described in Step 1 of the Streptavidin Flash Plate Assay above were performed.

Step 2:

30 The reaction was stopped by adding 20 μ l of 7.5M Guanidine Hydrochloride. 50 μ l of the stopped reaction was transferred to the Streptavidin filter plate (SAM[™] Biotin Capture Plate, Promega, catalog no. V7542) and the reaction was incubated on the filter for 1-2 minutes before applying vacuum.
35

The plate was then washed using a vacuum manifold as follows: 1) 4 x 200 μ l/well of 2M NaCl; 2) 6 x 200 μ l/well of 2M NaCl with 1% H_3PO_4 ; 3) 2 x 200 μ l/well of diH_2O ; and 4) 2 x 100 μ l/well of 95% Ethanol. The membranes were then allowed to air dry completely before adding scintillant.

5 The bottom of the plate was sealed with white backing tape, 30 μ l/well of Microscint 20 (Packard Instruments, catalog no. 6013621) was added. The top of the plate was sealed with clear sealing tape, and the plate then counted using the Packard TopCount with the appropriate settings for [^{33}P] with liquid scintillant.

10 Procedure for Phosphocellulose Filter Plate Assay:

Step 1:

The enzymatic reactions were performed as described in Step 1 of the Streptavidin

15 Flash Plate Assay (above) utilizing KKGGRARTSSFAEPG (SEQ.ID.NO.: 16) as the substrate in place of biotin-GGRARTSSFAEPG.

Step 2:

20 The reaction was stopped by adding 20 μ l of 0.75% H_3PO_4 . 50 μ l of stopped reaction was transferred to the filter plate (UNIFILTER™, Whatman P81 Strong Cation Exchanger, White Polystyrene 96 Well Plates, Polyfiltronics, catalog no. 7700-3312) and the reaction incubated on the filter for 1-2 minutes before applying vacuum.

The plate was then washed using a vacuum manifold as follows: 1) 9 x 200 μ l/well of 0.75% H_3PO_4 ; and 2) 2 x 200 μ l/well of diH_2O . The bottom of the plate was sealed with white backing tape, then 30 μ l/well of Microscint 20 was added. The top of the plate was sealed with clear sealing tape, and the plate counted using the Packard TopCount with the appropriate settings for [^{33}P] and liquid scintillant.

30 PKA Assay

Each individual PKA assay consists of the following components:

- 1) 10 μ l 5X PKA assay buffer (200 mM Tris pH7.5, 100 mM $MgCl_2$, 5mM 2-mercaptoethanol, 0.5 mM EDTA)
- 2) 10 μ l of a 50 μ M stock of Kemptide (Sigma) diluted into water
- 3) 10 μ l ^{33}P -ATP (prepared by diluting 1.0 μ l ^{33}P -ATP [10 mCi/ml] into 200 μ l of a 50 μ M stock of unlabeled ATP)
- 4) 10 μ l appropriate solvent control dilution or inhibitor dilution
- 5) 10 μ l of a 70 nM stock of PKA catalytic subunit (UBI catalog # 14-114) diluted in 0.5 mg/ml BSA.

The final assay concentrations were 40 mM Tris pH 7.5, 20 mM $MgCl_2$, 1 mM 2-mercaptoethanol, 0.1 mM EDTA, 10 μ M Kemptide, 10 μ M ^{33}P -ATP, 14 nM PKA and 0.1 mg/ml BSA.

Assays were assembled in 96 deep-well assay plates. Components #3 and #4 were premixed and in a separate tube, a mixture containing equal volumes of components #1, #2, and #5 was prepared. The assay reaction was initiated by adding 30 μ l of the components #1, #2, and #5 mixture to wells containing ^{33}P -ATP and inhibitor. The liquid in the assay wells was mixed and the assay reactions incubated for 20 minutes at room temperature. The reactions were stopped by adding 50 μ l 100 mM EDTA and 100 mM sodium pyrophosphate and mixing.

The enzyme reaction product (phosphorylated Kemptide) was quantitated using p81 phosphocellulose 96 well filter plates (Millipore). Each well of a p81 filter plate was filled with 75 mM phosphoric acid. The wells were aspirated and 170 μ l of 75 mM phosphoric acid was added to each well. A 30-40 μ l aliquot from each stopped PKA reaction was added to corresponding wells on the filter plate containing the phosphoric acid. The peptide was trapped on the filter following the application of a vacuum. The filters were washed 5X by filling wells with 75 mM phosphoric acid followed by aspiration. After the final wash, the filters were allowed to air dry. 30 μ l scintillation fluid was added to each well and the filters counted on a TopCount (Packard).

PKC Assay

Each PKC assay consists of the following components:

- 5 1) 5 μ l 10X PKC co-activation buffer (2.5 mM EGTA, 4mM CaCl_2)
- 2) 10 μ l 5X PKC activation buffer (1.6 mg/ml phosphatidylserine, 0.16
 mg/ml diacylglycerol, 100 mM Tris pH 7.5, 50 mM MgCl, 5 mM 2-
 mercaptoethanol)
- 10 3) 5 μ l ^{33}P -ATP (prepared by diluting 1.0 μ l ^{33}P -ATP [10 mCi/ml] into
 100 μ l of a 100 μ M stock of unlabeled ATP)
- 4) 10 μ l of a 350 μ g/ml stock of myelin basic protein (MBP, UBI) diluted
15 in water
- 5) 10 μ l appropriate solvent control or inhibitor dilution
- 6) 10 μ l of a 50ng/ml stock of PKC (mix of isoforms from UBI catalog #
20 14-115) diluted into 0.5 mg/ml BSA

Final assay concentrations were as follows: 0.25 mM EGTA, 0.4 mM
CaCl, 20 mM Tris pH 7.5, 10 mM MgCl, 1 mM 2-mercaptoethanol, 0.32 mg/ml
phosphatidylserine, 0.032 mg/ml diacylglycerol, 10 μ M ^{33}P -ATP, 70 μ g/ml MBP, 10
25 ng/ml PKC, 0.1 mg/ml BSA.

Assays are performed using 96 deep well assay plates. In each assay
well 10 μ l of solvent control or appropriate inhibitor dilution with 5 μ l ^{33}P -ATP
(components #5 and #3) were premixed. In a separate tube, a mixture containing
equal volumes of components #1, #2, #4, and #6 was prepared. The assay reaction
30 was initiated by adding 35 μ l of the components #1, #2, #4, and #6 mixture to wells
containing ^{33}P -ATP and inhibitor. The liquid in the assay wells was thoroughly mixed
and the assay reactions incubated for 20 minutes at room temperature. The reactions
were stopped by adding 100 mM EDTA (50 μ l) and 100 mM sodium pyrophosphate
(50 μ l) and mixing. Phosphorylated MBP was collected on PVDF membranes in 96
35 well filter plates and quantitated by scintillation counting.

The results from testing the compounds described in Examples 1-2 in the assays described above are shown in Table 1:

TABLE 1

	GSK3 Peptide Substrate IC ₅₀ (μM)				Counter screens IC ₅₀ (μM)	
	Akt-1	Akt-1 delta PH	Akt2	Akt3	PKA	PKC
Compound 2	3.88	>50	>50	>50	>40	>40
Compound 1	10.5	>50	>50	>50	>40	>40

EXAMPLE 7

Cell based Assays to Determine Inhibition of Akt/PKB

Cells (for example LnCaP or a PTEN^(-/-) tumor cell line with activated Akt/PKB) were plated in 100mm dishes. When the cells were approximately 70 to 80% confluent, the cells were refed with 5mls of fresh media and the test compound added in solution. Controls included untreated cells, vehicle treated cells and cells treated with either LY294002 (Sigma) or wortmanin (Sigma) at 20 μM or 200 nM, respectively. The cells were incubated for 2 hrs, and the media removed. The cells were washed with PBS, scraped and transferred to a centrifuge tube. They were pelleted and washed again with PBS. Finally, the cell pellet was resuspended in lysis buffer (20 mM Tris pH8, 140 mM NaCl, 2 mM EDTA, 1% Triton, 1 mM Na Pyrophosphate, 10 mM β-Glycerol Phosphate, 10 mM NaF, 0.5 mM NaVO₄, 1 μM Microcystine, and 1x Protease Inhibitor Cocktail), placed on ice for 15 minutes and gently vortexed to lyse the cells. The lysate was spun in a Beckman tabletop ultra centrifuge at 100,000 x g at 4°C for 20min. The supernatant protein was quantitated by a standard Bradford protocol (BioRad) and stored at -70°C until needed.

Proteins were immunoprecipitated (IP) from cleared lysates as follows:
For Akt1/PKB α , lysates are mixed with Santa Cruz sc-7126 (D-17) in NETN
(100mM NaCl, 20mM Tris pH 8.0, 1mM EDTA, 0.5% NP-40) and Protein A/G
Agarose (Santa Cruz sc-2003) was added. For Akt2/PKB β , lysates were mixed in
5 NETN with anti-Akt-2 agarose (Upstate Biotechnology #16-174) and for Akt3/PKB γ ,
lysates were mixed in NETN with anti-Akt-3 agarose (Upstate Biotechnology #16-
175). The IPs were incubated overnight at 4° C, washed and separated by SDS-
PAGE.

Western blots were used to analyze total Akt, pThr308 Akt, pSer473
10 Akt, and downstream targets of Akt using specific antibodies (Cell Signaling
Technology): Anti-Total Akt (cat. no. 9272), Anti-Phospho Akt Serine 473 (cat. no.
9271), and Anti-Phospho Akt Threonine 308 (cat. no. 9275). After incubating with
the appropriate primary antibody diluted in PBS + 0.5% non-fat dry milk (NFDm) at
4°C overnight, blots were washed, incubated with Horseradish peroxidase (HRP)-
15 tagged secondary antibody in PBS + 0.5% NFDm for 1 hour at room temperature.
Proteins were detected with ECL Reagents (Amersham/Pharmacia Biotech
RPN2134).

EXAMPLE 8

20

Heregulin Stimulated Akt Activation

MCF7 cells (a human breast cancer line that is PTEN^{+/+}) were plated at
1x10⁶ cells per 100mm plate. When the cells were 70 – 80% confluent, they were
refed with 5 ml of serum free media and incubated overnight. The following
25 morning, compound was added and the cells were incubated for 1 – 2 hours, heregulin
was added (to induce the activation of Akt) for 30 minutes and the cells were analyzed
as described above.

EXAMPLE 9

30

Inhibition Of Tumor Growth


In vivo efficacy of an inhibitor of the growth of cancer cells may be
confirmed by several protocols well known in the art.

Human tumor cell lines which exhibit a deregulation of the PI3K
35 pathway (such as LnCaP, PC3, C33a, OVCAR-3, MDA-MB-468 or the like) are

injected subcutaneously into the left flank of 8-12 week old female nude mice (Harlan) on day 0. The mice are randomly assigned to a vehicle, compound or combination treatment group. Daily subcutaneous administration begins on day 1 and continues for the duration of the experiment. Alternatively, the inhibitor test
5 compound may be administered by a continuous infusion pump. Compound, compound combination or vehicle is delivered in a total volume of 0.1 ml. Tumors are excised and weighed when all of the vehicle-treated animals exhibited lesions of 0.5 - 1.0 cm in diameter, typically 4 to 5.5 weeks after the cells were injected. The
10 average weight of the tumors in each treatment group for each cell line is calculated.

WHAT IS CLAIMED IS:

1. A compound of the formula A:



A

wherein

R^1 represents phenyl, furyl, thienyl or pyridinyl, any of which groups may be optionally substituted with one, two or three substituents, independently selected

from:

- a) halogen;
- b) C_{1-4} alkyl;
- c) C_{1-4} alkoxy;
- d) cyano;
- e) di(C_{1-4} alkyl)amino;
- f) hydroxy;

R^2 represents amino- C_{1-6} alkyl, C_{1-4} alkylamino-(C_{1-6})alkyl, di(C_{1-4} alkyl)amino-(C_{1-6})alkyl, hydroxy-(C_{1-6})alkyl or C_{1-4} alkoxy-(C_{1-6})alkyl, any of which groups may be optionally substituted;


R^3 represents hydrogen or C_{1-6} alkyl; and

R^4 independently represents hydrogen, C_{1-6} -alkyl, halogen, HO- or C_{1-6} alkyl-O;

r is 0, 1 or 2;

or a pharmaceutically acceptable salt or stereoisomer thereof.

2. The compound according to Claim 1 of the formula A-I:



A-I

wherein

R^2 is as defined with reference to formula A above;

R^4 is selected from: C_{3-7} cycloalkyl and phenyl, any of which groups may be optionally substituted;

r is 0, 1 or 2;

s is 0, 1, 2 or 3; and

R^5 independently represents halogen, C_{1-4} alkyl or C_{1-6} alkoxy;

or a pharmaceutically acceptable salt or stereoisomer thereof.

3. The compound according to Claim 1 which is:

2,2,*N,N*-tetramethyl-*N*-(3-phenyl-[1,2,4]triazolo[3,4-*a*]phthalazin-6-yl)-propane-1,3-diamine

N'-[3-(4-Methoxy-phenyl)-[1,2,4]triazolo[4,3-*a*]phthalazin-6-yl]-2,2,*N,N*-tetramethyl-propane-1,3-diamine

or a pharmaceutically acceptable salt thereof.

4. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 1.

5. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 3.

5 6. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 4.

10 7. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 5.

15 8. A pharmaceutical composition made by combining the compound of Claim 1 and a pharmaceutically acceptable carrier.

9. A process for making a pharmaceutical composition comprising combining a compound of Claim 1 and a pharmaceutically acceptable carrier.

AMENDED CLAIMS

[Received by the International Bureau on 06 September 2002 (06.09.02);
original claims 6-9 amended; new claims 10-15 added; remaining claims
unchanged]

5. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 3.

6. A method of selectively inhibiting one or more of the isoforms of Akt in a mammal which comprises administering to the mammal a therapeutically effective amount of a compound of Claim 1.

7. A method of selectively inhibiting one or more of the isoforms of Akt in a mammal which comprises administering to the mammal a therapeutically effective amount of a compound of Claim 3.

8. A method of selectively inhibiting one or more of the isoforms of Akt in a mammal which comprises administering to the mammal a therapeutically effective amount of a composition of Claim 4.

9. A method of selectively inhibiting one or more of the isoforms of Akt in a mammal which comprises administering to the mammal a therapeutically effective amount of a composition of Claim 5.

10. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 1.

11. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 3.

12. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 4.

13. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a composition of Claim 5.

AMENDED SHEET (ARTICLE 19)

14. A pharmaceutical composition made by combining the compound of Claim 1 and a pharmaceutically acceptable carrier.

15. A process for making a pharmaceutical composition comprising combining a compound of Claim 1 and a pharmaceutically acceptable carrier.

SEQUENCE LISTING

<110> Stanley F. Barnett
Andrew Pate Owens

<120> INHIBITORS OF AKT ACTIVITY

<130> 20838

<150> 60/282,824

<151> 2001-04-10

<160> 16

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Completely Synthetic Amino Acid Sequence

<400> 1

Cys Thr Gly Cys Gly Gly Cys Cys Gly Cys
1 5 10

<210> 2

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> Completely Synthetic Amino Acid Sequence

<400> 2

Gly Thr Ala Cys Gly Cys Gly Gly Cys Cys Gly Cys Ala Gly
1 5 10

<210> 3

<211> 40

<212> PRT

<213> Artificial Sequence

<220>

<223> Completely Synthetic Amino Acid Sequence

<400> 3

Cys Gly Cys Gly Ala Ala Thr Thr Cys Ala Gly Ala Thr Cys Thr Ala
1 5 10 15
Cys Cys Ala Ser Thr Glu Ala Gly Cys Gly Ala Cys Gly Thr Gly Gly
20 25 30
Cys Thr Ala Thr Thr Gly Thr Gly
35 40

<210> 4

<211> 33
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Completely Synthetic Amino Acid Sequence

<400> 4
 Cys Gly Cys Thr Cys Thr Ala Gly Ala Gly Gly Ala Thr Cys Cys Thr
 1 5 10 15
 Cys Ala Gly Gly Cys Cys Gly Thr Gly Cys Thr Gly Cys Thr Gly Gly
 20 25 30
 Cys

<210> 5
 <211> 24
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Completely Synthetic Amino Acid Sequence

<400> 5
 Gly Thr Ala Cys Gly Ala Thr Gly Cys Thr Gly Ala Ala Cys Gly Ala
 1 5 10 15
 Thr Ala Thr Cys Thr Thr Cys Gly
 20

<210> 6
 <211> 45
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Completely Synthetic Amino Acid Sequence

<400> 6
 Gly Ala Ala Thr Ala Cys Ala Thr Gly Cys Cys Gly Ala Thr Gly Gly
 1 5 10 15
 Ala Ala Ala Gly Cys Gly Ala Cys Gly Gly Gly Gly Cys Thr Gly Ala
 20 25 30
 Ala Gly Ala Gly Ala Thr Gly Gly Ala Gly Gly Thr Gly
 35 40 45

<210> 7
 <211> 45
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Completely Synthetic Amino Acid Sequence

<400> 7
 Cys Cys Cys Cys Thr Cys Cys Ala Thr Cys Thr Cys Thr Thr Cys Ala
 1 5 10 15
 Gly Cys Cys Cys Cys Gly Thr Cys Gly Cys Thr Thr Thr Cys Cys Ala
 20 25 30
 Thr Cys Gly Gly Cys Ala Thr Gly Thr Ala Thr Thr Cys

35 40 45

<210> 8
 <211> 36
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Completely Synthetic Amino Acid Sequence

<400> 8
 Gly Ala Ala Thr Thr Cys Ala Gly Ala Thr Cys Thr Ala Cys Cys Ala
 1 5 10 15
 Thr Gly Ala Gly Cys Gly Ala Thr Gly Thr Thr Ala Cys Cys Ala Thr
 20 25 30
 Thr Gly Thr Gly
 35

<210> 9
 <211> 30
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Completely Synthetic Amino Acid Sequence

<400> 9
 Thr Cys Thr Ala Gly Ala Thr Cys Thr Thr Ala Thr Thr Cys Thr Cys
 1 5 10 15
 Gly Thr Cys Cys Ala Cys Thr Thr Gly Cys Ala Gly Ala Gly
 20 25 30

<210> 10
 <211> 48
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Completely Synthetic Amino Acid Sequence

<400> 10
 Gly Gly Thr Ala Cys Cys Ala Thr Gly Gly Ala Ala Thr Ala Cys Ala
 1 5 10 15
 Thr Gly Cys Cys Gly Ala Thr Gly Gly Ala Ala Ala Gly Cys Gly Ala
 20 25 30
 Thr Gly Thr Thr Ala Cys Cys Ala Thr Thr Gly Thr Gly Ala Ala Gly
 35 40 45

<210> 11
 <211> 33
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Completely Synthetic Amino Acid Sequence

<400> 11
 Ala Ala Gly Cys Thr Thr Ala Gly Ala Thr Cys Thr Ala Cys Cys Ala
 1 5 10 15

Thr Gly Ala Ala Thr Gly Ala Gly Gly Thr Gly Thr Cys Thr Gly Thr
 20 25 30
 Cys

<210> 12

<211> 30

<212> PRT

<213> Artificial Sequence

<220>

<223> Completely Synthetic Amino Acid Sequence

<400> 12

Gly Ala Ala Thr Thr Cys Gly Gly Ala Thr Cys Cys Thr Cys Ala Cys
 1 5 10 15
 Thr Cys Gly Cys Gly Gly Ala Thr Gly Cys Thr Gly Gly Cys
 20 25 30

<210> 13

<211> 49

<212> PRT

<213> Artificial Sequence

<220>

<223> Completely Synthetic Amino Acid Sequence

<400> 13

Gly Gly Thr Ala Cys Cys Ala Thr Gly Gly Ala Ala Thr Ala Cys Ala
 1 5 10 15
 Thr Gly Cys Cys Gly Ala Thr Gly Gly Ala Ala Ala Thr Gly Ala
 20 25 30
 Gly Gly Thr Gly Thr Cys Thr Gly Thr Cys Ala Thr Cys Ala Ala Ala
 35 40 45
 Gly

<210> 14

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Completely Synthetic Amino Acid Sequence

<400> 14

Glu Tyr Met Pro Met Glu
 1 5

<210> 15

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Completely Synthetic Amino Acid Sequence

<400> 15

Gly Gly Arg Ala Arg Thr Ser Ser Phe Ala Glu Pro Gly

1. 5. 10

<210> 16

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Completely Synthetic Amino Acid Sequence

<400> 16

Lys Lys Gly Gly Arg Ala Arg Thr Ser Ser Phe Ala Glu Pro Gly

1

5

10

15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US02/10881

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/5025; C07D 487/04

US CL : 514/248; 544/234

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/248; 544/234

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE STRUCTURE SEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4,788,186 A (OCCELLI et al.) 29 November 1988, see especially columns 1-3 and Table III, examples 29,31 and 32.	1,2,4,8,9

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 JUNE 2002

Date of mailing of the international search report

09 JUL 2002

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

EMILY BERNHARDT

Telephone No. (703) 308-1235

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.